

# Gravitational Magnetic Force on the Star S2 Orbiting the Galactic Center Supermassive Black Hole Leading to Dark Matter Force

Delso J

Bachelor's Degree in Physics by Zaragoza University, Spain

**Corresponding Author:** Jesus Delso Lapuerta, Bachelor's Degree in Physics by Zaragoza University, Spain.  
E-mail: [jesus.delso@gmail.com](mailto:jesus.delso@gmail.com)

**Received:** 📅 May 23, 2021; **Accepted:** 📅 May 29, 2021; **Published:** 📅 June 08, 2021

## Abstract

Similarly to the spin-orbit interaction there is a magnetic field acting on the star S2 induced by the supermassive black hole gravitational field. The gravitational magnetic force is derived from Special Relativity force transformations. Energy associated with this new force is stored in the dark matter energy tensor  $D(4, 0)$  related to the Riemann curvature.

Dark matter gravity is generated by  $D(4, 0)$  energy related directly to the  $S(4, 0)$  tensor, however this gravity is attributed to exotic particles never detected, galaxies in our universe are rotating with such speed that the gravity generated by their observable matter could not possibly hold them together.

Total Energy  $T(4, 0)$  and energy tensors are denied to complete the General Relativity field equations. The Ricci decomposition is a way of breaking up the Riemann curvature tensor into three orthogonal tensors,  $Z(4, 0)$ , Weyl tensor  $C(4, 0)$  and  $S(4, 0)$ .

**Keywords:** *Dark Matter; Expansion Of The Universe; Riemann Curvature; Ricci Decomposition; Special Relativity; Gravitational Magnetic Field; Gravitational Magnetic Force; Star S2; Galactic Center Supermassive Black Hole; General Relativity Field Equations*

## Gravitational Magnetic Force on the Star S2 Orbiting the Galactic Center Supermassive Black Hole

In quantum physics, in the spin-orbit interaction there is a magnetic field  $B$  acting on the electron in the rest frame of the electron, where  $v$  is the velocity of the electron and  $E$  is the electric field, in the non-relativistic limit,  $\gamma^{-1} = (1 - \beta^2)^{1/2}$ ,  $\beta = v/c$ ,  $c$  the speed of light,  $\gamma \approx 1$  [11].

$$B = -\frac{v \times E}{c^2} \quad (1)$$

A particle without electric charge ( $q = 0$ ) and rest mass  $m_0$  is orbiting a gravitational source with rest mass  $M_0$ ,  $m_0 \ll M_0$  and velocity  $v$  at its closest approach distance  $r$ , since  $q = 0$  the only force is the gravitational force.

The gravitational magnetic field is derived from Special Relativity force transformations [12], when velocities point to the same direction a repulsive gravitational magnetic force is induced  $F_{Bg}$ .

$$F_g = G \frac{M_0 m_0}{r^2}, F_{Bg} = \gamma^3 (-2\beta^2 + \beta^4) F_g \quad (2)$$

When velocities point to the opposite direction an attractive gravitational magnetic force is induced  $F_{Bg}$ .

$$F_g = G \frac{M_0 m_0}{r^2}, F_{Bg} = \gamma^3 (+2\beta^2 + \beta^4) F_g \quad (3)$$

So, with  $v = 7650$  km/s,  $r = 120$  AU,  $M_0 = 4.31$  million  $M_0$ ,  $m_0 = 10$   $M_0$ , [13] we obtain the  $F_{Bg}$  value on the star S2 orbiting the

Galactic center supermassive black hole at its closest approach

$$F_{Bg} = (0.00130399869725077405480262194772) F_g \quad (4)$$

$$F_{Bg} = 46048607207718957314308202785.368 \text{ Nw} \quad (5)$$

$$F_{Bg} (m_0^{-1} \gamma^{-1}) = 0.0023144098 \text{ m/s}^2 \quad (6)$$

$F_{Bg}$  is an extra attractive gravitational magnetic force pulling from the star; however this force is attributed to exotic particles never detected. You can also see discussions about dark matter and the expansion of the Universe in [10].

**Conformal energy  $U$  defined as a combination of  $C$  and the Hodge dual of  $C$ , dark matter energy  $D$  defined as a combination of  $S$  and the Hodge dual of  $S$ , similar definitions for  $V/Z$  and  $T/R$**

The Ricci decomposition is a way of breaking up the Riemann curvature tensor into three orthogonal tensors,  $Z$ , Weyl tensor  $C$  and  $S$ ,  $S$  tensor generates the dark matter gravity [6-9].

$$R_{ijkl} = Z_{ijkl} + C_{ijkl} + S_{ijkl} \quad (7)$$

$$S_{ijkl} = \frac{1}{12} R (g_{il} g_{jk} - g_{ik} g_{jl}) \quad (8)$$

$$Y_{jk} = R_{jk} - \frac{1}{4} R g_{jk}, Z_{ijkl} = \frac{1}{2} (Y_{il} g_{jk} - Y_{jl} g_{ik} - Y_{ik} g_{jl} + Y_{jk} g_{il}) \quad (9)$$

where  $R_{abcd}$  is the Riemann tensor,  $R_{ab}$  is the Ricci tensor,  $R$  is

the Ricci scalar (the scalar curvature).

The conformal energy tensor  $\mathbf{U}$  can be defined as a combination of  $\mathbf{C}$  and the Hodge dual of  $\mathbf{C}$  [1-3].

$$U_{abcd} = 1/8\pi(C_{amcd}C_{bcd}^m + *C_{amcd} * C_{bcd}^m + C_{abcn}C_{abd}^n + *C_{abcn} * C_{abd}^n) \quad (10)$$

The new dark matter energy tensor  $\mathbf{D}$  can be defined as a combination of  $\mathbf{S}$  and the Hodge dual of  $\mathbf{S}$

$$D_{abcd} = 1/8\pi(S_{amcd}S_{bcd}^m + *S_{amcd} * S_{bcd}^m + S_{abcn}S_{abd}^n + *S_{abcn} * S_{abd}^n) \quad (11)$$

The new energy tensor  $\mathbf{V}$  can be defined as a combination of  $\mathbf{Z}$  and the Hodge dual of  $\mathbf{Z}$

$$V_{abcd} = 1/8\pi(Z_{amcd}Z_{bcd}^m + *Z_{amcd} * Z_{bcd}^m + Z_{abcn}Z_{abd}^n + *Z_{abcn} * Z_{abd}^n) \quad (12)$$

The new Total Energy tensor  $\mathbf{T}$  can be defined as a combination of the Riemann tensor  $\mathbf{R}$  and the Hodge dual of  $\mathbf{R}$

$$T_{abcd} = 1/8\pi(R_{amcd}R_{bcd}^m + *R_{amcd} * R_{bcd}^m + R_{abcn}R_{abd}^n + *R_{abcn} * R_{abd}^n) \quad (13)$$

## Hodge Dual Definitions

The Hodge dual definition for Electromagnetic tensor and Weyl tensor [4]

$$*F_{ab} = \frac{1}{2}\epsilon_{abln}F^{ln} \quad (14)$$

$$*C_{abcd} = \frac{1}{2}\epsilon_{abln}C_{cd}^{ln} \quad (15)$$

The Hodge dual definition for dark matter  $\mathbf{S}$  tensor,  $\mathbf{Z}$  and  $\mathbf{R}$  tensors

$$*S_{abcd} = \frac{1}{2}\epsilon_{abln}S_{cd}^{ln}, *Z_{abcd} = \frac{1}{2}\epsilon_{abln}Z_{cd}^{ln}, *R_{abcd} = \frac{1}{2}\epsilon_{abln}R_{cd}^{ln} \quad (16)$$

Weyl tensor  $\mathbf{C}$  (4, 0) is related to the new Conformal Energy tensor  $\mathbf{U}$  (4,0). Dark matter tensor  $\mathbf{S}$  (4, 0) is related to the new dark matter energy tensor  $\mathbf{D}$  (4, 0).  $\mathbf{Z}$  (4, 0) tensor is related to the new energy tensor  $\mathbf{V}$  (4, 0). Riemann tensor  $\mathbf{R}$  (4, 0) is related to the new Total Energy tensor  $\mathbf{T}$  (4, 0).

## Complete General Relativity Field Equations

$$Z_{ab} - \frac{1}{2}Zg_{ab} + \Lambda_z g_{ab} = -k_z V_{ab} \quad (17)$$

$$C_{ab} - \frac{1}{2}Cg_{ab} + \Lambda_c g_{ab} = -k_c U_{ab} \quad (18)$$

$$S_{ab} - \frac{1}{2}Sg_{ab} + \Lambda_s g_{ab} = -k_s D_{ab} \quad (19)$$

$$R = Z + C + S \quad (20)$$

$$\kappa T = k_z V + k_c U + k_s D \quad (21)$$

$$\Lambda = \Lambda_z + \Lambda_c + \Lambda_s \quad (22)$$

$$\kappa T_{abcd} = k_z V_{abcd} + k_c U_{abcd} + k_s D_{abcd} \quad (23)$$

In the general theory of relativity the Einstein field equations relate the geometry of spacetime to the distribution of matter [5].

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \Lambda g_{\mu\nu} = -\kappa T_{\mu\nu} \quad (24)$$

## Conclusions

The maximum acceleration on the star S2 due to the magnetic gravitational force is 0.0023144098 at its closest approach, the supermassive black hole gravitational field is inducing an extra attractive gravitational magnetic force pulling from the star, and however this force is attributed to exotic particles never detected.

When a charged particle is moving it generates a magnetic field that interacts with charged particles in motion, similarly when a particle with mass is moving it generates a gravitational magnetic field that interacts with other particles in motion.

This new force, the gravitational magnetic force has been detected by astronomers for decades but has been misinterpreted, in a galaxy there are no dark matter particles generating the dark matter force, moving stars generate gravitational magnetic fields that interact with other stars in motion.

The gravitational magnetic force is derived from Special Relativity force transformations, if Special Relativity is correct the gravitational magnetic force is irrefutable.

## References

1. Jesus Delso Lapuerta (2020) On the conformal energy tensor defined as a combination of Weyl tensor and the Hodge dual of Weyl tensor Vixra.org Relativity and Cosmology, 2-3.
2. Roger Penrose (2006) El camino a la realidad Random House Mondadori, *Barcelona* 619.
3. Roger Penrose (2006) El camino a la realidad Random House Mondadori, *Barcelona* 629.
4. Roger Penrose (2006) El camino a la realidad Random House Mondadori, *Barcelona* 1255.
5. Albert Einstein (1985) El significado de la relatividad Planeta-De Agostini, *Barcelona* 175.
6. Besse Arthur L (1987) Einstein manifolds, *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*, vol. 10, Berlin, New York: Springer-Verlag, pp. xii+510.
7. Sharpe RW (1997) *Differential Geometry: Cartan's Generalization of Klein's Erlangen Program*, Springer-Verlag, New York, ISBN 0-387-94732-9. Section 6.1 discusses the decomposition. Versions of the decomposition also enter into the discussion of conformal and projective geometries, in chapters 7 and 8.
8. Singer IM, Thorpe JA (1969) The curvature of 4-dimensional Einstein spaces, *Global Analysis (Papers in Honor of K. Kodaira)*, Univ. Tokyo Press 355–365.
9. Robert Resnick (1977) *Introducción a la Teoría Especial de la Relatividad* Editorial Limusa, *México DF* 139.
10. Delso J (2021) On Dark Matter Gravity Force Generated by the Expansion of the Universe. *OSP Journal of Physics and Astronomy 2: JPA-2-111*.
11. Robert Eisberg, Robert Resnick (1979) *Física Cuántica. Átomos, moléculas, sólidos, núcleos y partículas* Editorial Limusa, *México DF* 329.
12. Delso J (2021) On Gravitational Magnetic Field Derived from Special Relativity Leading to Dark Matter Force. *OSP Journal of Physics and Astronomy 2: JPA-2-115*.
13. Wikipedia (2021) S2 (star) [wikipedia.org](https://en.wikipedia.org/wiki/S2_(star)).